

# PHQ-9 Quick Depression Assessment

## For initial diagnosis:

1. Patient completes PHQ-9 Quick Depression Assessment in your presence
2. If there are at least 4 ✓s in the unshaded section (Scoring >1 for question 1-8, or >0 for question 9), consider a depressive disorder. Add score to determine severity.

### **Consider Major Depressive Disorder**

- if there are at least 5 ✓s in the unshaded section (one of which corresponds to Question 1 or 2)

### **Consider Other Depressive Disorder**

- if there are 2-4 ✓s in the unshaded section (one of which corresponds to Question 1 or 2)

**Note:** Since the questionnaire relies on patient self-report, all responses should be verified by the clinician, and a definitive diagnosis is made on clinical grounds taking into account how well the patient understood the questionnaire, as well as other relevant information from the patient. Diagnoses of Major Depressive Disorder or Other Depressive Disorder also require impairment of social, occupational, or other important areas of functioning (Question 10) and ruling out normal bereavement, a history of a Manic Episode (Bipolar Disorder), and a physical disorder, medication, or other drug as the biological cause of the depressive symptoms.

## To monitor severity over time for newly diagnosed patients or patients in current treatment for depression:

1. Patients may complete questionnaires at baseline and at regular intervals (eg, every 2 weeks) at home and bring them in at their next appointment for scoring or they may complete the questionnaire during each scheduled appointment.
2. Add up ✓s by column. For every ✓: "Several days" = 1 "More than half the days" = 2 "Nearly every day" = 3
3. Add together column scores to get a TOTAL score.

## Interpretation of Total Score

Total Score	Depression Severity
1-4	Minimal depression
5-9	Mild depression
10-14	Moderate depression
15-19	Moderately severe depression
20-27	Severe depression

*Reproduced with the permission of Pfizer Limited*